

3 (a) Two metal cuboids P and Q are in thermal contact with each other.

(i) P and Q are in thermal equilibrium.

State what is meant by the term thermal equilibrium.

.....  
 .....  
 ..... [2]

(ii) Data for P and Q are given in Table 3.1.

**Table 3.1**

	P	Q
specific heat capacity/ $\text{J kg}^{-1} \text{K}^{-1}$	390	910
mass/kg	0.54	0.37

P and Q are initially both at the same temperature.

P is supplied with 24 kJ of thermal energy. After some time, P and Q are once again both at the same temperature as each other.


P and Q are perfectly insulated from the surroundings.

Determine the change in temperature  $\Delta T$  of Q.

$\Delta T = \dots\dots\dots$  K [3]

(b) Nitrogen may be assumed to be an ideal gas. A fixed amount of nitrogen gas is contained at a constant pressure of  $1.6 \times 10^5$  Pa.

The variation of the volume  $V$  of the gas with the temperature  $\theta$  of the gas is shown in Fig. 3.1.



**Fig. 3.1**

(i) The temperature of the nitrogen gas is increased from  $0^\circ\text{C}$  to  $210^\circ\text{C}$ . Determine the work done on the gas.

work done =  $\dots\dots\dots$  J [3]

(ii) Determine the number  $N$  of molecules of nitrogen gas.

$N = \dots\dots\dots$  [2]

(iii) The mass of a nitrogen molecule is  $4.7 \times 10^{-26}$  kg.

Calculate the root-mean-square (r.m.s.) speed of a nitrogen molecule at  $210^\circ\text{C}$ .

r.m.s. speed =  $\dots\dots\dots$   $\text{ms}^{-1}$  [2]