


4. Adult male guppies (*Poecilia reticulata*) exhibit genetically determined spots, while juvenile and adult female guppies lack spots. In a study of selection, male and female guppies from genetically diverse populations were collected from different mountain streams and placed together in an isolated environment containing no predators.

The study population was maintained for several generations in the isolated area before being separated into two groups. One group was moved to an artificial pond containing a fish predator, while a second group was moved to an artificial pond containing no predators. The two groups went through several generations in their new environments. At different times during the experiment, the mean number of spots per adult male guppy was determined as shown in the figure below. Vertical bars in the figure represent two standard errors of the mean (SEM).



- (a) **Describe** the change in genetic variation in the population between 0 and 6 months and **provide** reasoning for your description based on the means and SEM.
- (b) **Propose** ONE type of mating behavior that could have resulted in the observed change in the number of spots per adult male guppy between 6 and 20 months in the absence of the predator.
- (c) **Propose** an evolutionary mechanism that explains the change in average number of spots between 6 and 20 months in the presence of the predator.
-
5. Genetically modified crops have been developed that produce a protein that makes the plants resistant to insect pests. Other genetic modifications make the crops more resistant to chemicals that kill plants (herbicides).
- (a) **Describe** TWO potential biological risks of large-scale cultivation and use of such genetically modified plants.
- (b) For each of the risks you described in part (a), **propose** a practical approach for reducing the risk.