



2. A student designs an experiment to study the reaction between  $\text{NaHCO}_3$  and  $\text{HC}_2\text{H}_3\text{O}_2$ . The reaction is represented by the equation above. The student places 2.24 g of  $\text{NaHCO}_3$  in a flask and adds 60.0 mL of 0.875 M  $\text{HC}_2\text{H}_3\text{O}_2$ . The student observes the formation of bubbles and that the flask gets cooler as the reaction proceeds.
- Identify the reaction represented above as an acid-base reaction, precipitation reaction, or redox reaction. Justify your answer.
  - Based on the information above, identify the limiting reactant. Justify your answer with calculations.
  - The student observes that the bubbling is rapid at the beginning of the reaction and gradually slows as the reaction continues. Explain this change in the reaction rate in terms of the collisions between reactant particles.
  - In thermodynamic terms, a reaction can be driven by enthalpy, entropy, or both.
    - Considering that the flask gets cooler as the reaction proceeds, what drives the chemical reaction between  $\text{NaHCO}_3(s)$  and  $\text{HC}_2\text{H}_3\text{O}_2(aq)$ ? Answer by drawing a circle around one of the choices below.
 

|               |              |                           |
|---------------|--------------|---------------------------|
| Enthalpy only | Entropy only | Both enthalpy and entropy |
|---------------|--------------|---------------------------|
    - Justify your selection in part (d)(i) in terms of  $\Delta G^\circ$ .
  - A student prepares a solution containing equimolar amounts of  $\text{HC}_2\text{H}_3\text{O}_2$  and  $\text{NaC}_2\text{H}_3\text{O}_2$ . The pH of the solution is measured to be 4.7. The student adds two drops of 3.0 M  $\text{HNO}_3(aq)$  and stirs the sample, observing that the pH remains at 4.7. Write a balanced, net-ionic equation for the reaction between  $\text{HNO}_3(aq)$  and the chemical species in the sample that is responsible for the pH remaining at 4.7.