


Question 3: Version J

The following information applies to all parts.

3. Nitrous acid, HNO_2 , is a weak acid that ionizes according to Equation 1.



Part A

Identify a conjugate acid-base pair in the equation. Be sure to clearly label which is the acid and which is the base.

Part B

A solution of $\text{HNO}_2(aq)$ with an initial concentration of 0.125 M has a pH of 2.09.

- i. **Calculate** the value of $[\text{H}_3\text{O}^+]$ in the solution. Show the work that leads to your answer.
- ii. **Calculate** $[\text{HNO}_2]$ at equilibrium. Show the work that leads to your answer.

Part C

In an experiment, the solution of 0.125 M $\text{HNO}_2(aq)$ is heated to 333 K. The new equilibrium concentrations are determined to be $[\text{HNO}_2] = 0.114 \text{ M}$, $[\text{NO}_2^-] = 0.0109 \text{ M}$, and $[\text{H}_3\text{O}^+] = 0.0109 \text{ M}$.

- i. **Calculate** the value of K_a for HNO_2 at 333 K. Show the work that leads to your answer.
- ii. Is the reaction represented by Equation 1 endothermic or exothermic? **Justify** your answer by comparing K_a values at 298 K and 333 K.


The following information applies to parts D, E, and F.

In a second experiment, a student is given a bottle containing HNO_2 with an unknown molarity. To determine the concentration of the $\text{HNO}_2(aq)$ in the bottle, the student titrates 35.0 mL of the $\text{HNO}_2(aq)$ with 0.16 M $\text{NaOH}(aq)$ at 298 K, which reacts according to Equation 2.



The results of the titration are summarized in Figure 1.

Figure 1



Part D

Using Figure 1, **identify** the pH at the equivalence point.

Part E

Using Figure 1, **calculate** the molarity $\text{HNO}_2(aq)$ of in the bottle. Show the work that leads to your answer.

Part F

Draw an X on Figure 1 to represent a point in the titration where $[\text{HNO}_2] > [\text{NO}_2^-]$ in the reaction mixture.

Part G

Equilibrium constants at 298 K are shown in Table 1 for the acid ionization reaction represented by Equation 1, K_a ; the autoionization of water, K_w ; and the neutralization reaction represented by Equation 2, K_2 .

Table 1

Chemical Equation	Equilibrium Constant
$\text{HNO}_2(aq) + \text{H}_2\text{O}(l) \rightleftharpoons \text{NO}_2^-(aq) + \text{H}_3\text{O}^+(aq)$	$K_a = 5.6 \times 10^{-4}$
$\text{H}_2\text{O}(l) + \text{H}_2\text{O}(l) \rightleftharpoons \text{H}_3\text{O}^+(aq) + \text{OH}^-(aq)$	$K_w = 1.0 \times 10^{-14}$
$\text{HNO}_2(aq) + \text{OH}^-(aq) \rightarrow \text{NO}_2^-(aq) + \text{H}_2\text{O}(l)$	$K_2 = ?$

Calculate the value of K_2 for Equation 2 at 298 K. Show the work that leads to your answer.

Part H

The student decides to repeat the experiment, this time using an indicator, and has access to the indicators in Table 2.

Table 2

Indicator Name	Acid Color	pH Range of Color Change	Base Color
Methyl orange	Red	3.1–4.4	Yellow
Thymol blue	Yellow	8.0–9.6	Blue
Clayton yellow	Yellow	12.2–13.2	Orange

The student conducts a second titration by adding two drops of methyl orange indicator to 35.0 mL of the $\text{HNO}_2(aq)$ solution and titrating the solution with 0.16 M $\text{NaOH}(aq)$ until a color change from red to yellow occurs.

The student claims that methyl orange was the best choice for the indicator. Do you agree or disagree? **Justify** your answer.