

1. Respond to parts A, B, C, D, E, F, G, H, and I, and all subparts.
- A. Based on the data in the graph, **identify** the number of Common Tern breeding pairs in 1995.
- B. Based on the data in the graph, **describe** the trend in the number of Common Tern breeding pairs from 1995 to 2008.
- C. A group of students hypothesized that sea level rise will lead to an increase in Common Tern populations. **Describe** one way that the data in the graph refute this hypothesis.
- D. **Explain** how climate change can lead to sea level rise.
- E. Wading birds are a group of large-bodied birds that inhabit coastal wetlands all over the world. Researchers monitored the number of wading bird species at four coastal locations experiencing rising sea levels over a 5-year period. The data are shown in the following table.

Sea Level Rise and Number of Wading Bird Species at Four Coastal Locations


Year	Increase in Sea Level (m)	Location A	Location B	Location C	Location D
1	0.05	1	5	3	4
2	0.08	2	4	3	6
3	0.10	6	7	5	5
4	0.12	5	7	7	7
5	0.15	7	8	9	8

- i. **Identify** a hypothesis that researchers are likely investigating in the wading bird study.
- ii. **Identify** the dependent variable in this study.
- F. One year after the completion of the study of wading bird species described in Part E, locations A and B experienced an increase in the concentration of PCBs, known to be a persistent organic pollutant, running off into the water.
- If the researchers repeated their study, **explain** how this environmental change could affect the results of the study.
- G. **Describe** a regulating ecosystem service that can be provided by wetlands.
- H. Since 1990, there has been an increase in urbanization around the coastal wetlands that wading bird communities inhabit. Most of these bird species eat fish and locate prey visually while walking through shallow water. An increase in sediment was discovered in a wetland located near an urban area.
- Explain** how increased sediment might lead to a change in the size of the wading bird populations.
- I. **Describe** an environmental problem, other than an increase in sediment, that can occur in wetlands that is associated with increased urbanization and development.

AP ENVIRONMENTAL SCIENCE 2023 – FREE-RESPONSE QUESTIONS

Serengeti National Park is located in Tanzania, Africa. The park is located in a valley along a tectonic plate boundary called the East African Rift. Figure 1 shows the East African Rift and other plate boundaries near the east coast of Africa.

Figure 1. Tectonic Plate Movement near East Coast of Africa



AP ENVIRONMENTAL SCIENCE 2023 – FREE-RESPONSE QUESTIONS

Serengeti National Park contains a variety of tropical habitats but is dominated by wideopen lands that include a variety of grass species with scattered trees and shrubs. Most grass growth occurs after the March through May rainy season. Many of the park’s herbivores, such as wildebeest and zebra, migrate through the park in an annual cycle following the growth of new grass (Figure 2).

Figure 2. Annual Migration of Wildebeest in Serengeti National Park

