




Cart 1 of mass m_1 is held at rest above the bottom of an incline. Cart 2 has mass m_2 , where $m_2 > m_1$, and is at rest at the bottom of the incline. At time $t = 0$, Cart 1 is released and then travels down the incline and smoothly transitions to the horizontal section. The center of mass of Cart 1 moves a vertical distance of H , as shown. At time t_C , Cart 1 reaches the bottom of the incline and immediately collides with and sticks to Cart 2. After the collision, the two-cart system moves with constant speed v . Frictional and rotational effects are negligible.

(a) During the collision, is the impulse on Cart 1 from Cart 2 greater than, less than, or equal to the magnitude of the impulse on Cart 2 from Cart 1 ?


___ Greater than ___ Less than ___ Equal to

Justify your answer.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 2** on this page.

(b) On the following axes, draw graphs of the magnitude of the momentum of each cart as a function of time t , before and after t_C . The collision occurs in a negligible amount of time. The grid lines on each graph are drawn to the same scale.




(c) Show that the velocity v of the two-cart system after the collision is given by the equation

$$v = \sqrt{2g} \left(\frac{m_1}{m_1 + m_2} \right) \sqrt{H}.$$

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 2** on this page.

(d) A group of students use the setup to perform an experiment. They measure the mass of Cart 1 to be $m_1 = 0.250$ kg. The mass of Cart 2 is unknown. The students perform several trials and in each trial, Cart 1 is released from a different height H and the final velocity of the two-cart system is measured. The students graph v as a function of $\sqrt{H}$, as shown below.



i. Draw a line that represents the best fit to the data points shown.

ii. Use the best-fit line to calculate the mass of Cart 2.

GO ON TO THE NEXT PAGE.

Continue your response to **QUESTION 2** on this page.

(e) After the experiment, the students use a balance to measure the mass of Cart 2 and find it to be less than what was determined in part (d). To explain this discrepancy, one of the students proposes that the mass of Cart 1 was incorrectly measured at the beginning of the experiment. The students measure the mass of Cart 1 again and record a new value, m_1' .

Should the students expect that m_1' will be greater than 0.250 kg, less than 0.250 kg, or equal to 0.250 kg ?

___ $m_1' > 0.250$ kg ___ $m_1' < 0.250$ kg ___ $m_1' = 0.250$ kg

Justify your answer.

GO ON TO THE NEXT PAGE.